

Original Article

The geological environments in the accumulation of heavy metals in soils of Pinar del Río

J. M. Febles-González^{1*}

A.Y. Martínez-Robaina²

N. M. B Amaral-Sobrinho³

J.M. Febles-Díaz¹

Everaldo Zonta³

¹Universidad de La Habana, Calle 16 no.114 e/1ra y 3ra, Miramar. Playa, La Habana, Cuba, CP 10300

²Facultad de Ciencias Forestales y Agropecuarias, Universidad de Pinar del Río, Calle Martí N° 270 F. Pinar del Río, Cuba, CP 10400

³Universidad Federal Rural de Rio de Janeiro, Seropédica Seropédica-Río de Janeiro, CEP: 23.897-000, Brasil

* Author for correspondence: febles@rect.uh.cu, jmifebles@cim.uh.cu

ABSTRACT

The research was carried out in the South Plain of the Pinar Río province with the objective of integrally evaluating the geological environments of soil formation and their influence on the processes erosion-sedimentation-contamination by heavy metals (HMs), in localities destined to tobacco crop. To this end, two units with different degrees of intervention were selected: the erosive surface (Consolation of the South unit) and the erosive - cumulative surface (units of Pinar Río, San Juan and Martinez and San Luis). The pedological prospecting work identified four main types of soils. They were at depths of 0-20 cm described and sampled, to characterize the main properties that influence the accumulation of HMs in soils. The analysis carried out indicated that natural conditions do not exert a marked influence on the accumulation and availability of HMs, where the reference values of Cd, Pb, Cu, Zn, Fe, Mn, Ni, Cr and Co are, lower than the standards of quality established for Cuban soils.

Key words: tobacco, pollution, soil formation, soil degradation

INTRODUCTION

Heavy metals (HMs) are naturally present in soils. The increase in its concentrations can occur due to natural processes or due to anthropogenic activities ^(1,2). The total natural concentration of HMs in soils depends, mainly, on the source material, the formation processes and the proportion of the components of the solid phase, this concentration is less evident in soils that originate from sediments ^(3,4).

The natural concentration of HMs in soils has two main origins: the weathering of the parent material in a tropical environment modulated by different manifestations of erosion (current and geological) and the degradation processes generated by human activity, especially agriculture ⁽⁵⁾. Increases in concentrations due to natural causes are determined by the chemical-mineralogical composition of the underlying stone material, as well as the dynamics of the pedogenesis-morphogenesis processes ^(6,7) where fixation by clay minerals and complexation by organic matter play a role a fundamental role ^(8,9) (Table 1).

Table 1. Ranges of heavy metal concentrations in the most abundant rock types

Heavy metals	Igneous rocks			Sedimentary rocks		
	Ultra basic	Basic	Granites	Limestone	Sandstone	Schist
Cd	0.12	0.13-0.2	0.09-0.20	0.028-0.10	0.05	0.20
Co	110-150	35-50	1	0,1-40	0,30	19-20
Cr	2000-2980	200	4	10-11	35	90-100
Cu	10-42	90-100	10-13	5.5-15	30	39-50
Hg	0.004	0.01-0.08	0.08	0.05-0.16	0.03-0.29	0.18-0.5
Mn	1040-1300	1500-2200	400-500	620-1100	4 – 60	850
Ni	2000	150	0.50	7-12	2-9	68-70
Pb	0.10-140	3-5	20-24	5.7-7	8-10	20-23
Sn	0.50	1-1.5	3-3.50	0.5-4	0.50	4-6
Zn	50-58	100	40-52	20-25	16-30	100-120

Source: ⁽⁹⁾

In this context, the genetic regularities and links of the sedimentation and contamination processes remain to be with the manifestations of soil erosion specified, whether of a geological nature in environments that are little disturbed or induced by agricultural activity ⁽¹⁰⁾. Taking into account the previous antecedents, this work has the objective of comprehensively evaluating the geological environments and their influence on the processes of erosion-sedimentation-contamination by HMs in the soils of the Southern Plain of Pinar del Río.

MATERIALS AND METHODS

The research was carried out in the Southern Plain of the Pinar del Río province (Figure 1). In the region known as “Macizo Tabacalero de Vuelta Abajo”, in which two representative units of the historical-natural formative environments were selected at the local level of the soils in little disturbed localities: the erosive surface (Consolación del Sur unit) and the erosive-cumulative surface (units of Pinar del Río, San Juan y Martínez and San Luís).




Figure 1. Spatial distribution of the soil sampling points in the Southern Plain of Pinar del Río, Cuba

Twelve main profiles were characterized in them and soil samples were taken in the upper, middle and lower sectors of the micro-relief flexures. This is in order to examine by means of the descriptive-comparative method at the depth of 0-20 cm, the dynamics, manifestation and intensity of morphogenetic processes, in correspondence with geological - geomorphological variations and agricultural use, which influence the accumulation of HMs

in soils (Table 2). For the chemical analysis and pseudo-total concentrations of HMs, 60 samples were collected in the same number of sampling points, in five groups of uncultivated soils in areas covered by spontaneous vegetation or forest fragments with a certain level of morphogenetic balance. The final sample of 1 kg of soil was obtained from 20 sub-samples taken in the form of a zig-zag; which were air dried, crushed and sieved to a diameter of 2 mm.

Table 2. Classification of the soils of the investigated sub-region of the Southern Plain of Pinar del Río, Cuba

N°	Extension superficial (ha)	Percentage of the total area	Genetic Classification Soils of Cuba ⁽¹¹⁾		Classification of Soils of Cuba ⁽¹²⁾	World Soil Reference Base ⁽¹³⁾
1	10 026	59.5	Leachate	Yellow	Leachate	Chromic acrysol,
			Quartzite Ferralitic		Yellowish Ferralitic	
2	2 608.2	5.12	Leachate	Reddish	ABA	Acrisol Aluminic
			yellow ferralitic	–	yellowish red	
3	1 609.2	15.5	Quartzite sandy		Arenosols	Arenosols
4	1 237.2	9.54	Alluvial		Fluvisol	Fluvisol
5	863.4	7.34	Bronw with carbonates		Bronw	Eutric Cambisol

Source: Own elaboration

The analyzes were carried out in laboratories of the Institute of Agronomy of the Federal Rural University of Rio de Janeiro, Brazil, between 2016 and 2018, according to criteria and guideline values of soil quality ⁽¹⁴⁾. The pH was determined with a potentiometer in relation to soil water 1: 2.5. The organic matter (OM) was by oxidation with potassium dichromate (0.2 mol L⁻¹) in an acid medium determined, the excess being valued with ammoniacal ferrous sulfate. Interchangeable calcium and magnesium were extracted using a 1-mol L⁻¹ KCl solution and the analysis was carried out by the complexometric method with EDTA. Extraction of assimilable phosphorus and exchangeable potassium was carried out using a Mehlich-1 solution (HCL 0.05 mol L⁻¹ and H₂SO₄ 0.025 mol L⁻¹). The reading was with a photocolimeter for phosphorus made and a flame photometer for potassium. The evaluations were according to the Rio de Janeiro State Subscriber and Liming Manual carried out ⁽¹⁵⁾. To determine pseudo-total concentrations of HMs, 1 g of the sieved soil sample was taken, digested by heating with the MARS Xpress[®] Digester, USEPA Method 3051A using inverted royal water ⁽¹⁶⁾. The resulting extracts were by EAA analyzed in a VARIAN-55B kit where the metallic elements Cd, Pb, Zn, Fe, Mn, Ni, Cu Cr and Co were

quantified. For statistical analyzes, the arithmetic mean and standard deviation were used; the means were compared using the ANOVA and Tukey tests ($p < 0.05$).

RESULTS AND DISCUSSION

Influence of geological-geomorphological conditions on the accumulation of heavy metals in the soils of the Southern Plain of Pinar del Río

The edaphic cover "archives" traits and properties inherited from past climatic and geological phases, which are not in balance with current edaphogenic processes ⁽¹⁷⁾, where the geological structure influences the origin of the MPs present today in the soils according to its zonal - spatial distribution (Table 3).

Table 3. Main rocks and mineralogical composition of the investigated regions

Formation	Lithology/diagnostic	Age	Thicknesses
Guane (gne N_2^2 Q_1^1)	Siliceous sands, sandy clays, gravels (angular and subangled), weakly cemented by clays with an indefinite lenticular layering.	Upper Pliocene - Lower Pleistocene	It can reach up to 50 m.
Guevara (gv Q_1^{1-2})	Plastic clays (montmorillonite and montmorillonite - kaolinitic), siliceous sands, fine gravels, fragments of ferritic shells (hardpan) with an indefinite, parallel stratification.	Lower-Middle Pleistocene.	It does not exceed 50 m.
Alluvial and marshy deposits (al Q_2 and p Q_2)	The alluvial deposits (al Q_2) are associated with the valleys and river terraces of rivers and streams, consisting mainly of quartziferous sands. Clay sands and sandy clays are also with intercalations of gravels of varied composition and dimensions documented. The marsh sediments (p Q_2) are only in the coastal zones distributed; clays and carbonaceous silts with vegetal remains represent them.		

Source: (18)

A significant characteristic of the soils is their transit to the basal material or underlying weathering crust, which is associated with the elution of the clays through the profile in an environment of greater stability (concave flexures of the micro slopes), confirmed dynamics in the morphologist - genetic observations made, corroborating what was stated ⁽¹⁹⁾. In some cases, these weathered sediments, either due to their location in the micro-relief or having a higher proportion of clays, are enriched with HMs, so that the soils can be classified as "contaminated" ⁽¹⁰⁾. However, these concentrations were found naturally or of low anthropic activity, due to the presence of these elements in the constituent minerals of the rocks and

elluvio-deluvial deposits characteristic of the geological formation environments, coinciding with studies carried out in similar regions ⁽³⁾.

Pedogenesis–morfogénesis dynamic and its influence in heavy metal accumulation

Through a comprehensive analysis, the current stage of morphogenesis is examined and, in a broader sense, the dynamics of some of the properties of soils that, due to their low content of organic matter and sandy texture ⁽²⁰⁾. Among other characteristics, they are very erodible ones, together with traditional management practices that have led to the progressive decline of their chemical, physical and biological fertility ⁽²¹⁾.

In this context, two sub-regions were from the interpretation of the landforms differentiated and in particular the pedogenesis-morphogenesis balance Northeast (erosive-cumulative surface) and South (cumulative surface). It is characterized by a different and complex paleogeographical evolution both the geological-geomorphological conditions, as well as by the conditions of use that at the same time have influenced the zonal - spatial distribution of the HMs (Figure 2). It presents variations in the edaphic cover, in which the alitic soils with low activity stand out red-yellowish clay (ABARA), Ferralitic Yellowish Leachate (FRAL) and Arenosols, among other types such as Fluvisol and Arenosol according to what was reported ⁽¹²⁾.





Figure 2. Soil distribution in the main geological-geomorphological surfaces in the Southern Plain of Pinar del Río, Cuba

Erosive - cumulative surface. Upper sector of Consolación del Sur unit

In it, the most accentuated slopes are located, both for the position it occupies in the context of the province and for the dissimilar processes that take place (Figure 3). They are classified as a scattering surface for creeping ⁽²²⁾, where the weathering-erosion processes have given rise to redeposited automorphic bark of the kaolinite-alitic-ferric type at a depth between 40-60 cm, associated with clay mantles, whose main morphological features conform to a lightly textured, generally acidic edaphic horizon Low humidification, low cation exchange capacity, desaturated with ferruginous concretions.



Automorphic surface. Transient - unstable

Ø=10 % Erosion due to areal runoff and trenches H=50-100 m

Vertical dismemberment = 2.5-25 m/km²

Horizontal dismemberment = 1 m/km²

Soil dominance ABARA

Natural vegetation: Marabou, aroma and grasslands

Deliver of porous sediments that descend towards areas of greater geomorphological stability

Semi-hydromorphic accumulation surface

Ø=8 % Erosion due to areal runoff and trenches H = 45-25 m

Vertical dismemberment = 2.5-25 m/km²

Horizontal dismemberment = 1 m/km²

Fluvisol soil prevalence

Natural vegetation: Marabou, aroma and grasslands

Clay-sandy sediment receptor in depresional zones

Figure 3. Distribution of the main soils in the Consolación del Sur Unit. Pinar del Río, Cuba

The development of the profile has been influenced by natural causes and anthropogenic factors, which have sometimes interrupted the evolutionary-sequential process described, leading to a combination of soils at the genetic level: Alitic with low red-yellowish clay

activity (ABARA), Ferralitic Yellowish Leachate (FRAL) and Fluvisol, where the pedological differentiation is highly contrasting. It is revealed by the changes in the chemical-mineralogical composition of the underlying stone material and the degree of evolution, attending to the topographic enclave they occupy.

Therefore, the ABARA and FRAL soil profiles are in a wavy relief distributed (Figure 3). The spatial configuration of the slope shapes are less stable ($\varnothing=8-10\%$; $H=50-100\text{ m}$). Fluvisol soils are basically confined to the flood valleys of rivers and flat sectors of the territory ($\varnothing=2-4\%$; $H=300-50\text{ m}$), with contributions of non-native materials and HPs in the upper thickness being common of the profiles that coincide with the results obtained ⁽²³⁾ in the Southern Plain of Pinar del Río.

In relation to physical properties, soils generally have a light texture, with a percentage of clay between 5-30 % at the level of the genetic horizons A+B0-50 cm. They are the most important for tobacco cultivation (Table 4), where 70–95 % of its mass is of sandy fractions made up, fundamentally kaolinite of good crystallization and in a lesser proportion illite and vermiculite, with traces of chlorite, being a regular increase in said minerals in depth according to the position they occupy in the micro-relief. This illuvial horizon eventually emerges on the surface when erosion is very strong.

Table 4. Physical properties of the main soils in little disturbed environments. Towns of Consolación del Sur municipality

Horizon/ Depth (cm)	Percentage of fractions in mm					Mg m ⁻³		Munsell Color
	A. Gross 2 - 0.2	A. Fine 0.2 - 0.02	Slime G. 0.02 - 0.01	Slime F. 0.01 - 0.002	C < 0.002	Density		
						From soil	Solid phase	
Profile C₁ (No apparent erosion)								
- ALYICAL LOW ACTIVITY CLAY RED-YELLOW -								
A₁ (0 - 12)	0.30	76.10	4.82	5.18	13.60	1.49	2.62	7.5YR 5/7
AB (12 - 33)	2.70	65.80	3.60	6.60	21.10	1.48	2.63	5YR 4/6
B₁ (33 - 45)	3.40	53.88	4.12	5.80	32.80	1.49	2.64	7.5 YR 6/8
(45 - 66)	1.60	49.40	5.30	5.90	37.90	1.50	2.66	2.5YR 5/7
B₂ (66 - 96)	0.10	46.42	5.72	7.53	38.87	1.52	2.67	2.5 YR 6/8
(96 - 129)	0.30	47.68	5.72	7.10	39.20	1.51	2.66	2.5 YR 6/8
BC (129 - 168)	0.10	40.50	7.20	7.22	47.00	1.52	2.67	2.5 YR 5/8
Profile C₂ (No apparent erosion)								
-- FERRALYTIC YELLOW LEACHATE --								
Ap (0- 20)	31.63	33.47	8.78	10.54	13.56	1.42	2.62	10 YR 6/4
AB (20 -42)	23.23	41.77	8.45	11.13	13.42	1.43	2.63	10 YR 7/5
B₁ (42 -76)	20.59	39.29	10.26	11.94	17.90	1.50	2.65	10 YR 7/8
B₂ (76 -95)	34.61	26.04	8.71	12.83	17.81	1.45	2.67	10 YR 6/8
BC (95 - 117)	44.59	23.63	7.77	12.69	19.30	1.48	2.67	10 YR 7/6
C (117 - 125)	48.36	25.45	6.54	12.46	21.19	1.50	2.70	7.8 YR 7/6
Profile C₅ (No apparent erosion)								
- FLUVISOL -								
A (0 -15)	11.02	81.78	0.80	3.50	3.90	1.40	2.67	10 YR6/8
AC (15 - 50)	9.72	74.28	11.10	2.40	2.50	1.42	2.67	10 YR7/9
C₁ (50 -90)	13.02	68.28	5.30	0.90	12.50	1.46	2.66	10 YR6/9
C₂ (90 - 120)	10.12	67.38	5.12	2.38	15.00	1.42	2.65	10 YR6/10
C₃ (120 - 200)	11.03	60.41	5.82	2.73	20.01	1.42	2.67	10 YR6/10

The analysis of the physical-chemical and chemical properties (Table 5), denotes an average value of organic matter of 3.44 %, qualifying on average according to ⁽¹⁵⁾, where the ABARA soil presents the lowest value as a manifestation of the development and maintenance of the aspects, degree of erodability and use indicated. The cation exchange capacity shows an average value of 15.69 cmol_c kg⁻¹ (low) in correspondence with the percentage of clay and the pH in the first horizons, together with the average contents of the exchangeable bases (mainly Ca and Mg) and sandy texture ⁽²³⁾.


Table 5. Physico-chemical and chemical properties of the soils in little disturbed environments in the Southern Plain of Pinar del Río, Cuba

Parameters	pH	Ca	Mg	K	CIC ¹	M.O ²	P ₂ O ₅	Clay
	(H ₂ O)	----- cmol.kg ⁻¹ -----				%	Mg kg ⁻¹	%
FRAL (n = 24). Consolación del Sur–San Juan y Martínez units								
Average	6.1	6.83	1.85	0.74	15.68	3.07	95.24	5.9
ABARA (n = 12). Consolación del Sur–San Juan y Martínez units								
Average	7.36	8.2	1.67	0.6	13.38	2.85	45.43	5.3
ARENOSOLS (n = 6). Unidad San Luís								
Average	7.33	10.8	2.8	0.73	17.09	3.22	57.99	4.6
FLUVISOL (n = 18). Consolación del Sur Unit								
Average	6.99	10.93	1.52	0.76	15.1	4.63	37.48	9.1
Average (n=60)	6.95	9.19	1.96	0.71	15.31	3.44	59.04	6.23
St Dev..	0.59	2.01	0.58	0.07	1.54	0.81	25.57	1.99

¹Exchange cation exchange capacity, ² organic matter, Std. Std - standard deviation

Cumulative area. Middle-lower sector Pinar del Río, San Juan y Martínez and San Luís units

This surface shows differences in soil properties mainly determined by the spatial configuration of the slopes at the meso and micro relief levels (Figure 4), which from a geological point of view are associated with the Guane + Guevara Formation ⁽²⁴⁾.



Automorphic surface. Transient - unstable

Provides sediment. May present covered karst

Ø= 10 % Areal runoff erosion creep H=30-50 m

Vertical dismemberment = 2.5-5 m/km²

Horizontal dismemberment = 1 m/km²

Soil dominance ABARA

Natural vegetation: Marabou, aroma and grasslands, which acts as a geological barrier

Semi-hydromorphic accumulation surface

Accumulation of sandy-clay sediments

Ø=2-6 % Erosion due to areal runoff and sedimentation in flood valleys H=20-30 m

Vertical dismemberment = <1 m/km²

Horizontal dismemberment: 0.5 m/km²

Predominance of soils Arenasols

Natural vegetation: Marabou, aroma and grasslands

Clay-sandy sediments (undifferentiated)

Figure 4. Distribution of the main soils. Pinar del Río, San Juan y Martínez and San Luís Units, Cuba

It covers the units (SJM and PR), with a hypsometry between H=30-50 m; where the sediments are characterized by their morphological and textural similarities, but with different ages given the joint action of the processes of erosion - transport - sedimentation, with interrelated functions that represent a geomorphological barrier. It has led to a differentiation of synthesis and decomposition of organic matter, promoting the formation of the ABARA, FRAL and Arenosols soils with a marked contrast and in close relationship with the lithogeomorphological conditions, a dynamic that corresponds to those studied for some authors ⁽²⁵⁾.

On the other hand, the ABARA and FRAL soils of the lower-middle sector occupy the largest area (Table 6), where the forming materials are a consequence of the redeposition of weathered eluvia containing certain amounts of mica, from the quartzite shale of the highest regions. While the Arenosols are at heights between 20-40 m distributed and they are characterized by a pedological differentiation of the cover layer at the meso relief level in the river flood valleys (interfluvial spaces 250-450 m) that they intercept the territory.

Table 6. Physical properties of the main soils in little disturbed environments. Municipalities of Pinar del Río, San Juan y Martínez and San Luís

Horizon/ depth (cm)	Percentage of fractions in mm				Mg m ⁻³		Munsell Color
	A. Gross 2 - 0.2	A. Fine 0.2 - 0.02	Slime 0.02 - 0.002	Clay < 0.002	From soil	Solid Phase	
Profile C₁ (No apparent erosion)							
- LOW ACTIVITY ALITHITIC CLAY RED-YELLOW-(SJM)							
A ₁₍₀₋₁₅₎	10.08	62.29	13.33	13.31			7.5YR 5/7
AB ₍₁₅₋₄₅₎	31.57	37.26	7.59	23.62			5YR 4/6
B ₁₍₄₅₋₉₀₎	5.90	42.12	7.83	44.15			7.5 YR 6/8
B ₂₍₉₀₋₁₅₀₎	4.05	35.07	8.72	52.16			2.5 YR 6/8
Profile C₂ (No apparent erosion)							
- LEACHATE YELLOW FERRALITICAL-(PR)							
A ₍₀₋₁₁₎	1.30	75.10	9.00	12.60		2.60	10 YR 6/4
A ₂₍₁₁₋₂₈₎	2.60	65.90	10.40	21.10	1.50		9.5YR 7/5
B ₁₍₂₈₋₄₄₎	3.45	53.44	10.01	33.00	1.66	2.66	9 YR 7/8
(44-66)	1.50	49.39	11.10	38.00			10 YR 6/8
B ₂₍₆₆₋₉₆₎	0.11	48.39	12.30	38.60	1.67	2.67	10 YR 7/6
(96-129)	0.29	48.8	12.11	39.17			
BC ₁₂₉₋₁₆₉	-----	39.5	15.50	45.00	-----	-----	7.8 YR 7/6
Profile C₅ (No apparent erosion)							
- ARENOSOLS-(SL)							
A ₁₍₀₋₁₄₎	0.40	84.6	5.50	9.50	1.54	2.55	10 YR6/8
A ₂₍₁₄₋₃₅₎	0.50	88.1	4.40	7.00	1.59	2.60	10 YR7/9
B ₁₍₃₅₋₅₂₎	0.40	86.9	4.20	8.50	1.60	2.60	10 YR6/9
B ₂₍₅₂₋₉₀₎	1.90	72.3	6.40	19.40	1.60	2.63	10 YR6/10
C ₍₉₀₋₁₈₀₎	1.90	72.4	3.10	22.60	1.65	2.65	10 YR6/10

In depth, a reduction in the diameter of the pores is manifested as it becomes more compact, caused by the accumulations of the finest fractions (<0.002 mm) in the profile that occupy the poral space from 50 cm depth (Table 6). It limits the penetration of the roots or does so by coating the aggregates, compressing, behavior reported in tobacco plots ⁽²⁶⁾.

Accumulation of heavy metals and their spatial distribution in the main soils of the Southern Plain of Pinar de Río

Regarding the origin of the main metallic elements, it has been strategic to consider the paleogeographic evolution of the Southern Plain of Pinar del Río, to know the origins of the current geochemical behavior of the main soils. A frequent feature is the abrupt transit of the B horizon to the old (ferrallitized) weathering crust, either *in situ* or redeposited, which regularly constitutes the C horizon of the soils ⁽²⁷⁾, generally of little biological activity and in most of the reclining cases between 120-150 cm deep.

The low natural concentrations of HMs (Table 7) are because these soils are supported mainly on transported materials of siliceous origin and kaolinized clay that do not retain a large amount of metal cations, due to the low content of clay and organic matter ⁽²⁸⁾. Since highly evolved soils formed under strong weathering processes present low natural concentrations of HMs, a criterion that coincides with similar investigations carried out ⁽²⁹⁾.

Table 7. Natural values of MPs in soils in little disturbed environments in the Southern Plain of Pinar del Río, Cuba

	Cd	Pb	Cu	Zn	Fe	Mn	Ni	Cr	Co
	----- mg kg ⁻¹ -----			----- g kg ⁻¹ -----		-----mg kg ⁻¹ -----			
FRAL (n = 24). Consolación del Sur – San Juan y Martínez Units									
Media	0.02 ^a	7.40cd	8.29a	26.72 ^a	9.86c	0.42ab	11.09ab	16.39b	7.06a
Desv. Est.	0.22	4.53	2.49	15.46	2.72	0.41	6.01	6.10	7.46
ABARA (n=12). Consolación del Sur – San Juan y Martínez Units									
Media	Nd	14.62bc	10.30 ^a	32.41 ^a	14.67bc	0.63a	15.15ab	22.46b	5.60a
Desv. Est.		3.19	1.48	3.12	2.41	0.15	1.50	1.94	0.81
ARENOSOLES (n = 6). San Luis Unit									
Media	Nd	1.72d	9.27 ^a	32.17 ^a	19.90b	0.69a	16.02a	19.46b	6.23a
Desv. Est.		2.67	5.31	14.45	6.40	0.20	3.31	24.41	1.42
FLUVISOL (n = 18). Consolación del Sur Unit									
Media	0.31 ^a	15.29b	12.06 ^a	40.36 ^a	15.22bc	0.34ab	12.72ab	27.22b	5.79a
Desv. Est.	0.41	0.28	1.48	5.14	3.17	0.11	2.56	3.11	1.00
¹CUBAN SOILS. (n = 33)									
Average	0.6	50	83	86	-	-	170	150	25

¹Average natural values of Cuban soils ⁽¹⁵⁾, Means with equal letters in the same column do not differ from each other. (Tukey; p <0.05)

Cr, Ni, Cu, Co and Zn are associated with primary minerals or co-precipitated with other secondary ones or strongly linked to Fe, Mn and Al oxides, they also accumulate in areas where depositions of soils removed by water erosion processes occur where metallic compounds are leached, a dynamic that corroborates the investigations carried out in similar regions ⁽³⁰⁾.

In the case of Cd, the concentrations are very low; only certain accumulations are in the FRAL, ABARA and Fluvisol soils observed, with values less than unity (Table 7). When comparing the concentrations of this element with the natural average values reported for Cuban soils ⁽³¹⁾, it is noted that the soils of the Consolación del Sur municipality have lower concentrations of this element. Cu and Zn presented the highest values in Fluvisol without differences with the rest of the soils, so it can be stated that the behavior of both elements is similar.

The Pb shows a heterogeneous behavior, in accordance with the Fe and Mn oxide contents given its strong capacity to adsorb this element. Something similar occurs with organic matter and high pH conditions that favor the formation of stable Pb chelates, resistant to leaching ⁽³²⁾. The FRAL soil has the lowest values of this element, a situation that is also expressed in its spatial distribution (Figure 5). The ABARA (14.62 mg kg⁻¹) and Fluvisol (15.29 mg kg⁻¹) soils present similar contents, in the Consolación del Sur - San Juan and Martínez units, respectively, due to the similarities between the characteristics of these two groupings in both surfaces.




Figure 5. Spatial behavior of Pb in the soils of the Southern Plain of Pinar del Río, Cuba

That is to say in ABARA soils due to the possible influence of the weathering crust of montmorillonite and montmorillonite-kaolinitic composition, which sometimes appears or appears close to the surface due to erosion. In Fluvisol soil, it is the result of accumulation of sediments with a sandy - clayey and clay - sandy texture with gravel intercalations of varied composition and dimensions, given their location in the depression areas as indicated. While Fe and Mn showed high and different values according to the type of soil (Figures 6 and 7), especially high concentrations in FRAL soils due to the nature of the source material and the formation process; where these elements are found mainly in the form of oxides, hydroxides or oxyhydroxides and are present in almost all soils ⁽³³⁾.




Figure 6. Spatial behavior of Fe in the soils of the Southern Plain of Pinar del Río, Cuba

Regarding their spatial distribution in the soils, the MPs Fe and Mn present similar behaviors in both sectors of the Southern Plain, as they are on materials supported with a similar chemical-mineralogical composition.




Figure 7. Spatial behavior of the Mn in the soils of the Southern Plain of Pinar del Río, Cuba

This geological-geomorphological conditioning offers a favorable environment for environmental equilibrium, if agro-ecological strategies for soil management are adopted. However, recultivation has been a common practice among farmers in the "Macizo Tabacalero Vizo Abajo", which consists of collecting sediments from the slopes near the rivers and applying it upstream. These sediments are generally enriched with PMs, toxic organic compounds, nutrients and organic matter, which could activate the retained toxic compounds, putting the quality of the productions and human health at risk, which is corroborated by what was reported in other regions of the country⁽³⁴⁾.

CONCLUSIONS

- The geological environments in the Southern Plain of Pinar de Río have exerted a notable influence on the composition, spatial distribution and accumulation of heavy metals, both in the current context and in the geographical palaeo, which generally show low natural concentrations compared to soils of other regions of the country.
- The results of this work constitute an approach to design more balanced management systems in risk scenarios of the tobacco massif of the Southern Plain of Pinar del Río and for others with similar edaphoclimatic and productive characteristics.

BIBLIOGRAPHY

1. Verma P, Agrawal M, Sagar R. Assessment of potential health risks due to heavy metals through vegetable consumption in a tropical area irrigated by treated wastewater. *Environment Systems and Decisions*. 2015;35(3):375–88.
2. Corrêa RM, Freire MBG dos S, Ferreira RLC, Freire FJ, Pessoa LGM, Miranda MA, et al. Atributos químicos de solos sob diferentes usos em perímetro irrigado no semiárido de Pernambuco. *Revista Brasileira de Ciência do Solo*. 2009;33(2):305–14.
3. Fadigas F de S, Sobrinho N, do Amaral MB, Mazur N, dos Anjos LH, Freixo AA. Proposição de valores de referência para a concentração natural de metais pesados em solos brasileiros. *Revista Brasileira de Engenharia Agrícola e Ambiental*. 2006;10(3):699–705.
4. Amundson R, Berhe AA, Hopmans JW, Olson C, Sztein AE, Sparks DL. Soil and human security in the 21st century. *Science*. 2015;348(6235):1261071.
5. Siddiqa A, Faisal M. Heavy Metals: Source, Toxicity Mechanisms, Health Effects, Nanotoxicology and Their Bioremediation. In: *Contaminants in Agriculture*. Springer; 2020. p. 117–41.
6. Wang F, Guan Q, Tian J, Lin J, Yang Y, Yang L, et al. Contamination characteristics, source apportionment, and health risk assessment of heavy metals in agricultural soil in the Hexi Corridor. *CATENA*. 2020;191:0341–8162.
7. Yang Y, Yang X, He M, Christakos G. Beyond mere pollution source identification: Determination of land covers emitting soil heavy metals by combining PCA/APCS, GeoDetector and GIS analysis. *Catena*. 2020;185:104297.
8. Sungur A, Vural A, Gundogdu A, Soylak M. Effect of antimonite mineralization area on heavy metal contents and geochemical fractions of agricultural soils in Gümüşhane Province, Turkey. *Catena*. 2020;184:0341–8162.
9. Peng S, Wang P, Peng L, Cheng T, Sun W, Shi Z. Predicting heavy metal partition equilibrium in soils: Roles of soil components and binding sites. *Soil Science Society of America Journal*. 2018;82(4):839–49.
10. Febles G, Amaral S, Pérez L, Zoffoli JH, Magalhães ML, Guedes N. Relation among the processes of erosion-sedimentation-pollution in soils from the Distrito Pecuário "Alturas de Nazareno", Cuba. *Cuban Journal of Agricultural Science*. 2014;48(2):173–9.

11. Hernández A, Pérez J, Bosch D, Rivero L, Camacho E, Ruiz J. Clasificación genética de los suelos de Cuba. Instituto de Suelos. Ministerio de la Agricultura. AGRINFOR. Ciudad de La Habana, Cuba. 1999;
12. Hernández JA, Pérez JJM, Bosch ID, Castro SN. Clasificación de los suelos de Cuba 2015. Mayabeque, Cuba: Ediciones INCA. 2015;93.
13. World reference base for soil resources 2014. International soil classification system for naming soils and creating legends for soil maps. Roma; 2015.
14. CONAMA-Conselho Nacional do Meio Ambiente Resolução n. 420 de 28 de dezembro de 2009. Dispõe sobre critérios e valores orientadores de qualidade do solo quanto à presença de substâncias químicas e estabelece diretrizes para o gerenciamento ambiental de áreas contaminadas por essas substâncias em decorrência de atividades antrópicas. publicada no Diário Oficial da União em Brasília, DF; 2009.
15. Freire LR. Manual de calagem e adubação do Estado do Rio de Janeiro. Embrapa Solos- Livro técnico (INFOTECA-E). 2013;
16. Technical Committee ISO/TC 190 S quality. Soil quality: Extraction of Trace Elements Soluble in Aqua Regia. International Organization for Standardization; 1995.
17. Alekseev AO, Kalinin PI, Alekseeva TV. Soil indicators of paleoenvironmental conditions in the south of the East European Plain in the Quaternary time. Eurasian Soil Science. 2019;52(4):349–58.
18. Peñalver-Hernández LL, Alonso JA, Rodríguez García A, Pérez Pupo R, Cruz Dávila F. Sobre la existencia de la Formación Camacho en la llanura sur de Pinar del Río. In: V Congreso Cubano de Geología y Minería, Memorias Geomin. 2003. p. 24–8.
19. Marrero A. Principales tipos de suelos de la cobertura de suelos de la llanura sur de Pinar del Río. [Tesis a Doctor en Ciencias Agrícolas]. [Habana]: Instituto de Suelos, Geografía de suelos; 1984.
20. Téllez OF, García JC, Obregón EF, García AT, Aguilar RL, Muñoz JM. Selección de alternativas en el tratamiento de suelos degradados utilizando métodos multicriterio. La Técnica. 2017;(17):6–17.
21. Hernández JML, Carcedo EC, de la Nuez Hernández E. Influencia de la agricultura de conservación sobre el suelo y el cultivo del tabaco en San Juan y Martínez, Cuba. Avances. 2015;17(4):318–26.

22. Loye A, Jaboyedoff M, Theule JI, Liébault F. Headwater sediment dynamics in a debris flow catchment constrained by high-resolution topographic surveys. *Earth Surface Dynamics*. 2016;4(2).
23. Robaina M, Yoán A, Febles González JM, do Amaral Sobrinho NM, Benítez Odio M, Morejón García M, et al. Alternancia de cultivos, su efecto sobre el suelo en zonas dedicadas a tabaco negro en Pinar del Río. *Centro Agrícola*. 2018;45(1):69–77.
24. Estévez Cruz E, Ordaz Hernández A, Hernández Santana JR. Deformaciones neotectónicas en el relieve fluvial de la Llanura Sur de Pinar del Río, Cuba. *Investigaciones geográficas*. 2017;(94).
25. Jaimez E, Ortega F. Paleorégimen hídrico en suelos con relictos edáficos. In: *Implicaciones en la reconstrucción paleoclimática del Pleistoceno superior para Cuba Occidental. IV Congreso de Geología y Minería (CD-Room. Textos/Geología del Cuaternario/210.html)*. La Habana. 2001.
26. Aroche EJA, Reina EM, Hernández JML. Diagnóstico inicial de la evolución de un suelo degradado. *Avances*. 2019;21(1):129–38.
27. Hernández Jiménez A. Pinar del Río-La Habana. Grabacion, Tour 4. La Habana ,Cuba; 1994.
28. Ramírez MG, Barrantes JAG, Thomas E, Miranda LAG, Pillaca M, Peramas LDT, et al. Heavy metals in alluvial gold mine spoils in the peruvian amazon. *Catena*. 2020;189.
29. do Amaral Sobrinho NMB, Chagas CI, Zonta E. Impactos ambientais provenientes da produção agrícola: experiências argentinas e brasileiras. Editora Autografia; 2016.
30. Zhang X, Yang H, Cui Z. Evaluation and analysis of soil migration and distribution characteristics of heavy metals in iron tailings. *Journal of Cleaner Production*. 2018;172:475–80.
31. Muñiz, O, Estévez, J., Quicute, S., Fraser, T, Vega, E., Montero, A. Contenido de metales pesados en áreas de agricultura intensiva de la Habana y Pinar del Río. [Internet]. 2020 Apr [cited 29/04/2020]. Available from: <https://www.google.es/search?hl=es&q=Mu%C3%B1iz,+O.,+Est%C3%A9vez,+J.,+Quicute,+S.,+Barrantes,+M.,+Fraser,+T.,+Vega,+E.,+Montero,+A.Contenido+de+metales+pesados+en+áreas+de+agricultura+intensiva+de+la+Habana+y+Pinar+del+Río+Informe+final+del+proyecto+07+perteneciente+PR-11:+Recursos+Naturales.,+>
32. Huang C, Zeng G, Huang D, Lai C, Xu P, Zhang C, et al. Effect of Phanerochaete chrysosporium inoculation on bacterial community and metal stabilization in lead-

- contaminated agricultural waste composting. *Bioresource technology*. 2017;243:294–303.
33. Alfaro MR, Montero A, Ugarte OM, do Nascimento CWA, de Aguiar Accioly AM, Biondi CM, et al. Background concentrations and reference values for heavy metals in soils of Cuba. *Environmental monitoring and assessment*. 2015;187(1):4198.
34. Ugarte OM, Alfaro MR, Álvarez AM, Álvarez JE, de Aguiar Accioly AM, Do Nascimento CWA. El Níquel en suelos y plantas de Cuba. *Cultivos Tropicales*. 2015;36:25–33.