



The reproduction of potato (*Solanum tuberosum* L.) tubers from their sexual seed

La reproducción de tubérculos de papa (*Solanum tuberosum* L.), a partir de su semilla sexual

Yadira López-Ramos*, Jorge L. Salomón-Díaz

Instituto Nacional de Ciencias Agrícolas (INCA), carretera San José-Tapaste, km 3½, Gaveta Postal 1, San José de las Lajas, Mayabeque, Cuba. CP 32 700.

ABSTRACT: Potato (*Solanum tuberosum* L.) is one of the most important food crops worldwide, due to its high nutritional values, its great demand among the population and its high production in a short time. For its propagation, propagules or tubers are used, which in Cuba are imported and they are very expensive; therefore, the objective of this review was to update the art state of potato tuber propagation from its sexual seed at international and national level. The review of the international bibliography showed that the research carried out has been aimed at obtaining new varieties, within the specialty of genetics and, in Cuba, the works carried out three decades ago were not concluded, so that an alternative technology has not yet been achieved for farmers to have access to the use of this technique. There is also no information available to show what remains to be done to achieve this goal. The results of the review allowed visualizing the benefits and limitations of this productive modality, for its possible economic, ecological and social benefits, if its implementation were achieved in Cuba, something that is not investigated at present.

Key words: food sovereignty, sustainability, technology, propagation, tubers, resilience.

RESUMEN: La papa (*Solanum tuberosum* L.) es uno de los cultivos alimenticios de mayor importancia a escala mundial, por sus altos valores nutricionales, su gran demanda entre los pobladores y por su elevada producción en corto tiempo. Para su reproducción se utilizan propágulos o tubérculos que para Cuba son importados y resultan muy costosos; por ello, el objetivo de esta revisión, fue lograr una actualización sobre el estado del arte de la reproducción de tubérculos de papa, a partir de su semilla sexual a escala internacional y nacional. La revisión de la bibliografía internacional mostró que las investigaciones realizadas han estado dirigidas a la obtención de nuevas variedades, dentro de la especialidad de la genética y, en Cuba, los trabajos realizados hace tres décadas, no fueron concluidos, por lo que no se ha logrado aún una tecnología alternativa al acceso de los agricultores para el uso de ésta técnica. Tampoco se cuenta con la información que muestre lo que falta por hacer para tal logro. Los resultados de la revisión permitieron visualizar las bondades y las limitaciones de esta modalidad productiva, para sus posibles beneficios económicos, ecológicos y sociales, si se lograra su implementación en Cuba, algo sobre lo cual no se investiga en la actualidad.

Palabras clave: soberanía alimentaria, sostenibilidad, tecnología, propagación, tubérculos, resiliencia.

INTRODUCTION

Potato is today one of the most important crops in the world food diet, ranking fourth after rice (*Oryza sativa* L.), wheat (*Triticum* spp.) and maize (*Zea mays* L.) (1). It has an annual production of 321.9 million tons on 18.6 million hectares, for an average yield of 17.3 t ha⁻¹. China is the world's leading potato producer with 73.7 million tons, followed by the Russian Federation with 36.4 t ha⁻¹, India with 25.0 t ha⁻¹ and Ukraine with 19.3 t ha⁻¹ (2).

According to the Ministry of Agriculture and Irrigation in Latin America, Peru remains the main potato producer in the region with an annual production of 5.3 million tons registered in 2019, whose volume serves to mitigate the food needs of that country, which has a population of 32 million consumers. This production volume of the Andean tuber, confirms Peru with the 14th place, among potato producers worldwide, only surpassed in descending order by Canada, Turkey, Iran, France, Belarus, Netherlands, Germany, Ukraine, United States, Poland, India, Russia and China, which occupies the first place (3).

*Author for correspondence: yadiral@inca.edu.cu

Received: 22/10/2020

Accepted: 20/09/2021



In Cuba, potato ranks first in production, among root and tuber crops, and an average of 7 515 ha are planted annually, with an average yield of 22.02 t ha⁻¹ and an annual production of 165 508.6 t (4).

All the potato produced worldwide is made through the use of tubers as planting material, from which Cuba is not exempt, with the disadvantage that such planting material is imported, mainly from Europe, together with a technological package for its production at too high a cost. This reality and considering the need to reduce imports to a minimum (5), it is necessary to find other more sustainable and resilient alternatives, which accompany the established technologies and, at the same time, contribute to increase its cultivation surface, with productive profitability towards a greater approach to Food Sovereignty (6). The reproduction of potato seed, through sexual seed, is an ancestral technique, derived from the first settlers of South America, the center of origin of this solanaceae (7).

However, the growing food demands on a global scale have caused the abandonment of traditional techniques, replaced by those that generate greater productivity over time, due to the growth of large cities, which demand large volumes of food in a short time and the demands of quality and uniformity imposed by the dynamics of international trade.

Potato, in particular, is a food consumed by more than one billion consumers worldwide, according to estimates, which has generalized propagation by propagules, which seems to be the most productive method in less time (2).

Cuba, which has scarce financial resources, imports at a high cost, all the propagules for the annual plantations, without the existence of a reproductive alternative to avoid imports or to contribute to supply the national demand. The reproduction of tubers through botanical seeds is an option that could be more sustainable and resilient for Cuba, to increase the production of this demanded food. For these reasons, the objective of this review was to achieve an update on the state of the art of potato tuber reproduction, from its sexual seed at international and national scale; this will allow visualizing its benefits and limitations and the possible economic, ecological and social benefits with its implementation in Cuba.

IMPORTANCE OF POTATO REPRODUCTION THROUGH BOTANICAL SEED

The literature reviewed indicates the existence of other planting alternatives through tubers, such as the use of buds that sprout from the tuber after its dormancy period, which allows a considerable increase in the number of plants per tuber. The production of seedlings through tissue culture, a novel method of agricultural biotechnology, and the production of tubers from botanical seed, which is included in the literature as an option whose main use has been in works for the propagation of varieties of genetic interest. The analysis of this review focuses on this last option (8).

Potato can be reproduced through its sexual seed, also known as botanical or true seed and by its acronym in English

as TPS (True Potato Seeds) (5,6). To achieve this, seeds are extracted from the berries and they are then used as reproduction material, an innovative technological option of low cost, since tubers obtained are free of many viruses and diseases and, even when there is no economic study of its costs, it is considered that this technique is relatively cheap, it does not require large refrigerated capacities for its conservation and the material resulting from its obtaining process is much easier to transport (9).

Sexual seed potato is currently used by farmers in at least a dozen countries, including China, India, Nepal, Bangladesh, Vietnam, Peru, Nicaragua, and Venezuela. Estimates of areas cultivated with sexual seed potato and planting materials derived from it are difficult to acquire because estimates depend on the extent to which subsequent generations of sexual seed potato materials can be traced and included in the estimates (10).

But the greatest importance of potato production from botanical seed is that it could be used by producers in situ and dispense with imported seed, which, moreover, it is very expensive. Although the commercialization of the product obtained may not be very attractive for marketing, due to its diversity of colors and shapes, it is still an important alternative for feeding families and neighbors in rural communities, not to mention its importance for animal feed (11).

RETROSPECTIVE ANALYSIS OF POTATO BREEDING TECHNOLOGY FROM SEXUAL SEED ON AN INTERNATIONAL SCALE

The literature reviewed makes reference to the fact that the potato was domesticated by the ancestors of Andean farmers and it was cultivated at least 7000 years ago. According to the distribution of the first known cultivated potatoes and the most related wild species, it seems likely that the cultivation of this tuber began in the Lake Titicaca region, between 3,500 and 4,500 meters above sea level. It is also in this region where there is the greatest variation in its cultivated forms and corresponding wild species (2).

With the passage of time, Andean man obtained hundreds of varieties, extending potato cultivation throughout almost the entire Andean region, occupying the high regions of Colombia, Ecuador, Peru, Bolivia and Chile. This period coincided with the arrival of the Spaniards in South America, who introduced it in Europe at the end of the 16th century, and it was subsequently dispersed throughout the world, due to commercial exchange (12).

Historically, it has been widely discussed about the species that gave rise to the potato cultivated between 6000 and 100,000 years ago, in the Andes of southern Peru.

Taking into account morphological and phytogeographical features, it is believed that *S. stenotomum* originated from the wild species *Solanum bukasovii*, *S. canasense* and *S. multisectum*, belonging to the *S. brevicaulis* complex, which is considered the first domesticated potato. This, in turn, would have given rise to *S. andigena*, through repeated processes of sexual polyploidization in different cultivation

areas, with the consequent interspecific and intervarietal hybridization, which allowed expanding the diversity and genetic adaptability of the Andean potato. Evidence from Andean folkloric history shows that highland farmers used sexual seed to periodically clean their planting materials and increase production (13).

POTATO REPRODUCTION THROUGH SEXUAL SEED IN MODERN TIMES

The first research on sexual seed potato of the modern era was carried out in the People's Republic of China, in 1959, and already in the beginning of 1960, started to produce on a large scale with this type of seed; therefore, this country is considered a pioneer in these studies (14,15).

In Latin America, it was the International Potato Center (CIP, according to its acronyms in Spanish) that in 1977 initiated a study on genetic improvement and agronomic experiments that became known as the technology for the reproduction of SSP. The first presentations on the subject were published in 1983-1984 in the Proceedings of Potato Research for the Year 2000 (16) and at the 28th Planning Conference on Innovative Methods in CIP for Potato Propagation (17).

An intense program of agronomic research was carried out by CIP's Department of Physiology at its headquarters and experimental stations located in Peru during the 1980's. From the agronomic point of view, the starting point was the experience developed by growing SSP seedlings produced from crosses. But in particular, reports of the innovative technology of extensive use of sexual seed potato (SSP) reported in China increased the enthusiasm in CIP circles. Thus, by 1984, more than 34 countries were experimenting with SSP and the results were used by farmers in five countries in the region (18).

There is also evidence of occasional use of botanical seed in Peru and Bolivia, such references to the use of this technique were described by the descendants of the Incas in the vicinity of Cuzco (19).

In 1989, Burton in his book "The Potato" in its (third edition), had integrated a chapter on propagation by botanical seed and since then, the technology has been experimented, integrated and adapted to a wide range of countries. In the 1990s, CIP reduced its research related to SSP, until it was phased out in the first decade of the 21st century, the 2000s. By this time, sufficient knowledge was available to formulate protocols for dormancy breaking and seed quality testing (10).

The positive experiences of experimentation with SSP in Central Asia have made it clear that this technology maintains the attraction of new countries and seed producing companies, which is reflected in the requests for courses on the subject, especially among researchers from Latin American countries.

However, not all options for improving the technology are fully explored. The technology can be further improved, especially through research related to genetics and seed quality (12).

SEXUAL REPRODUCTION OF POTATO IN CUBA. ITS IMPORTANCE

The oldest reference found on the sexual reproduction of potato in Cuba goes back to the results of the crossings carried out in the decade of the 80's of the XX century, with the purpose of selecting clonal varieties. The literature mentions that through the botanical seed, most of the viruses are not transmitted through it (10), except: the virus from the available material.

In the 90's of the 20th century, a research project was initiated on this subject called potato production from sexual seed. The experiments were conducted at the National Institute of Agricultural Sciences and at the Liliانا Dimitrova experimental station and the extension of results, in Isle of Youth special municipality. Aspects such as the selection of hybrid progenesis, fertilization and seed management were evaluated. Some of these results were reflected in the Guide for the production of potato seed tubers from sexual seed (12). At the end of the project, these studies were not continued. At present, there is a genetic improvement project and technology for obtaining botanical seed, which is being developed at INCA. However, there are still no conclusive results, reasons that advise to continue deepening in the subject, until achieving a technology on agroecological bases that makes possible its use in the productive practice, which would contribute to the Food Security of the country in times of scarce financial resources and would respond to the call of Cuban Science to work for the reduction of imports.

In Cuba, the production of tubers to be used as "seed" represents between 40-70 % of the cost of production. The high populations of aphids and other insects, in addition to the presence of diseases throughout the year, make it difficult to develop a national seed potato program, so it is imported every year with its technological package (20).

On the other hand, if it were possible to produce potato from botanical seed, abundant enough to supply needs of the communities, through the participation of local farmers, the importation of tubers as "seed" would gradually decrease, making the potato production process more sustainable and robust, in favor of the country's food sovereignty, even though the purpose of implementing an additional program is not intended to replace the established one.

ADVANTAGES AND LIMITATIONS OF POTATO PROPAGATION BY BOTANICAL SEED

Main advantages of potato propagation through botanical seed

When tuber production comes from botanical seed, genetic variability is rescued; plants obtained from botanical seed have tap roots, which can favor better anchorage and absorption of water and nutrients.

Sexual seed can be used for two different purposes: where potatoes can be produced directly in the field, from seed and in order to produce tubers to be used later as "seed" in a second harvest or even in repeated generations (10,25).

There are different sowing methods for tuber production: direct sowing in beds and transplanting of seedlings to beds (26).

Direct sowing in beds consists of sowing three or four seeds per nest at a distance of 10 x 10 cm; after emergence, the plants are thinned until 100/m² are left.

Transplanting of seedlings to beds consists of the following: the plants are grown in trays and transplanted to the beds 35 days after sowing, at a distance of 10 x 10 cm (this method requires more labor and delays the vegetative cycle of the plants by 20 days, due to the stress of transplanting).

When the two sowing methods were tested: direct sowing and transplanting (27), it was observed that the plants transplanted at 35 days suffered a delay of 10 days, compared to those of direct sowing and that more than 90 % of the tubers of direct sowing were formed between 45 and 55 days after sowing, while those of the transplanted plants were formed between 35 and 63 days later.

The total number of tubers at final harvest was similar in the two sowing methods, with a total duration of 110 days for direct sown plants and 125 days for transplanted plants.

The most commonly used technology is the production of tubers in beds.

When SSP is used in beds, there are fewer requirements for good germination and seedling growth. In addition, tuber seedling production does not require high transplanting inputs, plants do not suffer stress, and management is better controlled in small protected areas than in the field (23,28).

Density of population in beds for tubercle production

A high plant population has a positive effect on the number of usable tubers produced per unit area. Studies of stocking density in CIP show that the optimum plant population, after thinning, should be at least 100 plants/m²; to obtain this stocking density, botanical seed can be sown 10 cm apart between rows and two to three centimeters between seeds. Then the plants are thinned in each row, leaving more or less 10 cm between plants; thus a final distance of 10 x 10 cm between plants is obtained (21).

Sowing is done in points (nests) and the distance between each point should be 10 cm, obtaining 100 sowing points/m²-bed; in each point two or three seeds are deposited at a depth of 0.5 cm and then lightly covered with soil or substrate. After thinning and replanting in the points without plants, one plant should remain as a planting point. This density makes it possible to produce vigorous seedlings, increases air circulation and reduces the occurrence of fungal diseases, such as seedling drop and others that attack the foliage (10).

In experiments previously carried out by the author, testing two population densities of 50 and 100 plants/m², the highest number of total tubers was obtained with 100 plants/m² of different calibers in great number, except calibers greater

than 40 g, in which the number of tubers was equal to the number of plants (27).

Substratum preparation

Potato seedlings need a substrate containing a high percentage of organic matter and a very limited amount of clay. Too much clay hardens the substrate. A suitable substrate, measured in volumes, can be: four parts sand + four parts organic matter (compost or moss) and 0 to two parts loam or organic soil (21).

With the combined use of substrate in a 4:1 ratio of soil + cow manure, the use of biofertilizers in the co-inoculation of plants and a nitrogen pulse of 60 kg ha⁻¹, better results were obtained than when chemical fertilizers were used (29).

Some authors carried out studies where they tested three substrates: soil+cattle (2:1), soil+cattle manure (2:1) and soil+chemical fertilization, using two sowing methods (direct and by transplanting), obtaining satisfactory results with the substrate soil+cattle manure (2:1), both for the number of tubers per m², as well as for the fresh mass per m² (30).

When using different proportions of Ferrallitic Red soil compacted with cattle manure (2:1, 3:1 and 4:1), the best results were obtained with the 4:1 proportion, equivalent to 11.6 kg of manure/m² (86 t ha⁻¹) wet basis with 25% humidity (31).

Chemical and biological fertilization

Two weeks after germination (approximately 24 days after sowing), the first foliar fertilization is carried out, applying 10 liters of solution per 5 m² of beds, with a proportion of 1 gram of urea per liter of water. The second foliar fertilization is carried out six days after the first one, with a proportion of 2 grams of urea per liter of water (12).

The first fractional fertilization should be carried out 15 days after sowing, applying 275 g of urea or 390 g of ammonium nitrate; the second at 22 days after sowing, applying 623 g of urea or 917 g of ammonium nitrate and the third at 22 days after sowing, applying 623 g of urea or 917 g of ammonium nitrate and 780 g of potassium sulfate (25).

With the inoculation of potato sexual seeds with biofertilizers, such as *Pseudomonas fluorescens*, *Pseudomonas scepacia*, *Azospirillum brasilense* Sp-7 and the co-inoculation *Azospirillum brasilense* Sp-7+Mycorrhiza *Glomus fasciculatum*, sown in beds with soil+cow manure substrate at a ratio of 4: 1 and an application of 60 kg N ha⁻¹, it is possible to obtain higher yields in seed tubers per hectare than with traditional methods, with the consequent saving of chemical fertilizers and preserving the ecological balance of the ecosystem (29).

Hilling in flowerbeds

The maximum production of tubers per plant depends on a good application of soil around the plant to support it. If the earthing up is deficient, the stolons, when exposed to the air, do not become tubers but branches. Two hilling operations are carried out, the first immediately after thinning and the second 15 days later (24,25).

After the second hilling, the stems should be covered 4-6 cm above the initial level of the bed, the final conformation of the small furrows should be rounded in each row of plants and thus ensure that the roots and stolons do not come to the surface (12).

Hilling tends to increase the number of tubers per plant; it can also provide better protection against pests. Hilling can be affected by adding a layer of soil 4 to 6 cm high to the seedlings, dividing this amount into two hilling operations. The ridging should be completed before the plants touch each other (10,25).

Pests and diseases. Its management

Plant pathogens, such as nematodes, fungi, bacteria, mycoplasmas, viruses and viroids, can be transmitted from diseased plants to seed tubers. However, when propagation is by sexual seed potato these common potato diseases are not transmitted from diseased plants to sexual seed; therefore, they do not carry many of them. Only very few virus diseases and one viroid, so seed tubers from sexual seed potatoes have excellent phytosanitary quality, so they will not transmit many diseases to the next generation either. Well-documented reports of disease transmission by sexual seed potato are: potato T virus, tuber blight viroid, tobacco mottle virus, and Andean potato latent virus (32).

Harvest

Normally between 90-120 days after sowing, depending on the progeny used, plants will be ready to be harvested, which is recognized by a general yellowing. The foliage is then cut at ground level for harvesting 10-15 days later. Tubers that are exposed to the sun after defoliation should be covered with soil to prevent rotting. Light irrigations are applied during this stage to maintain soil moisture and fluffiness (10).

Seed preservation

Seed tubers from SSP should be stored under the same requirements of the Ramal Standard for potato crops, with the difference that sizes smaller than 30 mm should be stored in boxes of no more than 25 kg (12).

CONCLUSIONS

The SSP technology for potato production is a viable, pertinent and appropriate alternative that is adapted to the conditions of Cuban farmers for its possible implementation, which will undoubtedly contribute to local food security without the use of external resources and imported inputs.

RECOMMENDATIONS

1. That a program aimed at the production of sexual seed potatoes be initiated to put this alternative into production practice, with a view to increasing the availability of potato reproductive material for consumption and strengthening the availability of propagules for potato reproduction.

2. To develop, simultaneously, an investigation that defines the economic viability of the new proposal, assuming as a reference the expenses incurred for its acquisition, through its importation.
3. That research be initiated to solve problems related to genetic improvement and improvement of the phytotechnical management of potato cultivation during the process of tuber reproduction on agro-ecological bases.

BIBLIOGRAPHY

1. Izquierdo J. Manual técnico, producción de papa a partir de semilla sexual. Curso audiovisual, Juan Izquierdo ed. 1995; Available from: <https://biblioteca.inia.cl/handle/123456789/36191>
2. Estévez A. El cultivo de la papa en Cuba [Internet]. 2007 [cited 31/10/2021]. Available from: <http://www.ac-taf.co.cu/papa.html>
3. MIDAGRI. Perú se mantiene como primer productor de papa en América Latina [Internet]. [cited 31/10/2021]. Available from: <https://www.gob.pe/institucion/midagri/noticias/164182-peru-se-mantiene-como-primero-productor-de-papa-en-america-latina>
4. Martín R, Jeréz Mompí E. Evaluación del rendimiento en papa (*Solanum tuberosum*, L.) a partir del comportamiento de las temperaturas. *Cultivos tropicales* [Internet]. 2015;36(1):93-7. Available from: http://scielo.sld.cu/scielo.php?script=sci_arttext&pid=S0258-59362015000100012
5. Malagamba JP. Evaluación de Tecnología Agronómica para producción de Papa a partir de Semilla Botánica. 1983; Available from: https://pdf.usaid.gov/pdf_docs/PNABD714.pdf
6. Torres F. Guía de manejo de la semilla botánica (sexual) de papa para la producción de tubérculos- semilla. Programa Nacional de Papa, Ministerio de Agricultura y Ganadería, (MAG) Región I. "Las Segovias". Estelí, Nicaragua. 1991;
7. Franke G. *Nutzpflanzen der Tropen und Subtropen* [in 2 Bdn] [Internet]. Hirzel; 1980. Available from: <https://www.zvab.com/buch-suchen/titel/nutzpflanzen-der-tropen-und/autor/franke/>
8. Monteagudo H. *Principios de Agricultura*. Segunda edición modificada. a abana: Imprenta y Librería La propagandista; 1931. 451 (276-277).
9. Torres F, Olivas A. Producción de semilla sexual bajo las condiciones tropicales de Nicaragua. *Revista Latinoamericana de papa* [Internet]. 1993;5(1):1-19. Available from: https://books.google.es/books?hl=es&lr=&id=mV-kA4IH7UC&oi=fnd&pg=PA1&dq=Produccion%20de%20semilla%20sexual%20bajo%20las%20condiciones%20tropicales%20de%20Nicaragua&ots=ZBmGF96_3n&sig=WNNeMGMPHL_Tpn3gOnrIbFIDxmo#v=onepage&q=Produccion%20de%20semilla%20sexual%20bajo%20las%20condiciones%20tropicales%20de%20Nicaragua&f=false
10. Hernández A. Revisión bibliográfica La producción de papa por semilla sexual. *Cultivos Tropicales* [Internet]. 1999;20(4):57-71. Available from: <https://>

- www.researchgate.net/profile/Alejandro-F-Zardon/publication/321732050_Revisi3n_Bibliografica_La_produccion_de_papa_por_semilla_sexual/links/5a2ee7640f7e9bfe81700aa5/Revision-Bibliografica-La_produccion_de_papa_por_semilla_sexual.pdf
11. CIP. Procesamiento y usos de la papa [Internet]. International Potato Center. 2015 [cited 31/10/2021]. Available from: <https://cipotato.org/es/lapapa/procesamiento-y-usos-de-la-papa/>
 12. Zard3n AF, D3az JL, Mauri R, J.G C. Manual. Gu3a pr3ctica para la producci3n de tub3rculos- semilla de papa a partir de semilla sexual. [Internet]. 2015. Available from: https://www.researchgate.net/publication/321757460_Manual_Guia_practica_para_la_produccion_de_tuberculos-semilla_de_papa_a_partir_de_semilla_sexual
 13. Papa (CIP) CI de la. Informe Anual Centro Internacional de la Papa 1996 [Internet]. International Potato Center; 1997. 64 p. Available from: <https://books.google.com.co/books?id=6GULRRH0FMIC&printsec=frontcover#v=onepage&q&f=false>
 14. Bofu S, Yu QD, Vander Zaag P. True potato seed in China: past, present and future. American Potato Journal [Internet]. 1987;64(6):321-7. Available from: <https://link.springer.com/article/10.1007/BF02853524>
 15. Golmirzaie AM, Mendoza H. Estrategias de mejoramiento para la producci3n de semilla sexual de papa. CIP Circular. Centro Internacional de la Papa [Internet]. 1988;6(14). Available from: http://catalogo.infoagro.hn/cgi-bin/koha/opac-detail.pl?biblionumber=20314&shelfbrowse_itemnumber=20821
 16. Hooker W. Research for the potato in the year 2000 Lima Peru 22-27 February, 1982,. [Internet]. 1983. Available from: <https://kutuphane.tarimorman.gov.tr/vu-find/Record/10262/Description>
 17. Centro Internacional de Papa C. Innovative methods for propagating potatoes [Internet]. Lima, Per3; 1984. 333 p. Available from: https://pdf.usaid.gov/pdf_docs/PNABD687.pdf
 18. FAO. Organizaci3n de las Naciones Unidas para la Alimentaci3n y la Agricultura [Internet]. 2011. [cited 31/10/2021]. Available from: <https://www.fao.org/news/archive/news-by-date/2011/es/>
 19. Almekinders CJM, Chilver AS, Renia HM. Current status of the TPS technology in the world. Potato Research [Internet]. 1996;39(2):289-303. Available from: <https://link.springer.com/article/10.1007%252FBF02360921>
 20. Rodr3guez JM, Rodr3guez I, Serrano A, Salom3n JL, Gonz3lez E. Resultados y perspectivas de la producci3n de papa a partir de semilla sexual en la Rep3blica de Cuba. Evento Cient3fico: Producci3n de cultivos en condiciones tropicales. Res3menes.(1997: Quivic3n). 1997;13.
 21. Wiersema SG. Evaluaci3n de tecnolog3a para la producci3n de tub3rculos-semillas de semilla bot3nica de papa. Centro Internacional de la Papa; 1983.
 22. Graziano JE. Producir papa con semilla sexual. Una alternativa posible. Presencia [Internet]. 2011;22(56). Available from: <https://inta.gob.ar/sites/default/files/script-tmp-papasemilla.pdf>
 23. Almekinders CJM. On flowering and true potato seed production (*Solanum tuberosum*, L.). Wageningen Agricultural University. Wageningen, Netherlands [Internet]. 1995; Available from: <https://edepot.wur.nl/200067>
 24. Golmirzaie AM, Malagamba P, Pallais N. Breeding potatoes based on true seed propagation. Potato genetics. 1994;499-513.
 25. Pallais. N. Producci3n de semilla bot3nica de papa (TPS). 1986 [cited 31/10/2021]; Available from: <https://biblioteca.inia.cl/handle/123456789/64454>
 26. Wiersema SG. Produccion de tuberculos-semillas derivados de la semilla [Internet]. 1985. 13 p. Available from: <http://catalogo.infoagro.hn/cgi-bin/koha/opac-detail.pl?biblionumber=42190>
 27. Wiersema SG. A method of producing seed tubers from true potato seed. Potato Research [Internet]. 1986;29(2):225-37. Available from: <https://link.springer.com/article/10.1007%2FBF02357653>
 28. Struik PC, Wiersema SG. Seed potato technology [Internet]. Wageningen Academic Publishers; 1999. Available from: https://books.google.es/books?hl=es&lr=&id=TBPTDwAAQBAJ&oi=fnd&pg=PA355&dq=Seed+potato+technology&ots=I3TP6VAbDY&sig=eQcj8Mhan6_xghaFZDvwxC_DAM#v=onepage&q=Seed%20potato%20technology&f=false
 29. Hern3ndez A. Manejo agron3mico integral de sustratos, m3todos de siembra y biofertilizaci3n en la producci3n sostenible de tub3rculos-semilla de papa por semilla sexual. Cultivos tropicales [Internet]. 2001;22(2):21-7. Available from: <https://www.redalyc.org/pdf/1932/193215009003.pdf>
 30. Hern3ndez A, Corbera J, Plana R. Producci3n de tub3rculos semilla de papa mediante semilla bot3nica de polinizaci3n libre. Cultivos tropicales [Internet]. 1997;18(2):19-23. Available from: <https://ftp.inca.edu.cu/revista/1997/2/CT18205.pdf>
 31. Hernandez AF, Arzuaga J. Manejo agron3mico de sustratos, biofertilizantes y m3todos de siembra, en la producci3n sostenible de tuberculillos de papa por semilla sexual. 1998; Available from: <https://www.redalyc.org/pdf/1932/193215009003.pdf>
 32. Salazar LF. Potato viruses and their control [Internet]. International Potato Center; 1996. Available from: https://books.google.es/books?hl=es&lr=&id=1Nj4leyfu8C&oi=fnd&pg=PR9&dq=Potato+viruses+and+their+control&ots=-HY6fTtwfde&sig=rz_UvS_CvIOO7vAUvo_OitK34Uk#v=onepage&q=Potato%20viruses%20and%20their%20control&f=false